

DYNAMICAL NATURE OF THE NUCLEAR PSEUDOSPIN AND ITS ISOSPIN ASYMMETRY

R. LISBOA, M. MALHEIRO, A. DELFINO

Instituto de Física, Universidade Federal Fluminense, Niterói, Brazil

P. ALBERTO, M. FIOLHAIS

*Departamento de Física and Centro de Física Computacional, Universidade de
Coimbra, P-3004-516 Coimbra, Portugal*

M. CHIAPPARINI

Departamento de Física Teórica, Universidade do Estado do Rio de Janeiro, Brazil

Pseudospin symmetry in nuclei is investigated by solving the Dirac equation with Woods-Saxon scalar and vector radial potentials. We relate the pseudospin interaction with a pseudospin-orbit term in a Schroedinger-like equation for the lower component of the Dirac spinor. We show that this term gives a large contribution to the energy splittings of pseudospin partners, so that the near pseudospin degeneracy arises from a significant cancellation among the different terms in that equation. This is a manifestation of the dynamical character of this symmetry in the nucleus. We analyze the isospin dependence of the pseudospin symmetry in a nuclear isotope chain by including a vector-isovector potential V_ρ and a Coulomb potential and conclude that V_ρ gives the main contribution to the observed pseudospin isospin asymmetry.

1 Introduction

Pseudospin symmetry was introduced in the late 60's to account for a degeneracy in single-particle nucleon levels of certain nuclei, with quantum numbers $(n, \ell, j = \ell + 1/2)$ and $(n - 1, \ell + 2, j = \ell + 3/2)$ ^{1,2}. These levels have the same quantum number $\tilde{\ell} = \ell + 1$, called “pseudo” orbital angular momentum, and a “pseudo” spin quantum number, $\tilde{s} = 1/2$. In general, for a $n\ell j$ state, we have $\tilde{\ell} = \ell \pm 1$ if $j = \ell \pm 1/2$. Pseudospin symmetry is exact when doublets with $j = \tilde{\ell} \pm \tilde{s}$ are degenerate. Since then the origin of pseudospin symmetry in nuclei has been a subject of intense theoretical research.

Recently, Ginocchio ³ uncovered the relativistic character of the symmetry, noting that the pseudo-orbital angular momentum is the orbital angular momentum of the lower component of the Dirac spinor. This is an exact SU(2) symmetry for the Dirac Hamiltonian ⁴ with an attractive scalar potential, S , and a repulsive vector potential V , when they have the same magnitude, i.e., $\Sigma = S + V = 0$. This would explain why there are quasi-degenerate pseudospin doublets in nuclei, since in relativistic mean field (RMF) models nuclear sat-

uration arises from an extensive cancellation between a large attractive scalar potential and a large repulsive vector potential. However, since Σ acts as a binding potential, one cannot set $\Sigma = 0$ in the nucleus, because then there would not exist any bound states. Other authors have studied the role of pseudospin-orbit coupling in pseudospin symmetry ^{5,6}.

2 Dynamical pseudospin symmetry and isospin asymmetry

Pseudospin symmetry is related to an invariance under a SU(2) transformation of the Dirac Hamiltonian with central scalar S and vector V potentials

$$H = \boldsymbol{\alpha} \cdot \mathbf{p} + \beta(m + S) + V, \quad (1)$$

whose generators are ⁴ $S_i = s_i \frac{1}{2}(1 - \beta) + \tilde{s}_i \frac{1}{2}(1 + \beta)$ where $\tilde{s}_i = \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{p} s_i \frac{\boldsymbol{\sigma} \cdot \mathbf{p}}{p}$, $s_i = \sigma_i/2$. The commutator of S_i with (1) is zero if $\Sigma = 0$ or $\frac{d\Sigma}{dr} = 0$ ⁷.

The Dirac equation $H\Psi = \epsilon\Psi$ can be written as a pair of second-order differential equations for the upper and lower components. Setting $\Delta = V - S$, V and S being radial functions, and defining $\epsilon = E + m$, we have

$$p^2\Psi_+ = \frac{\Delta'}{E + 2m - \Delta} \left(\frac{\partial}{\partial r} - \frac{1}{r} \boldsymbol{\sigma} \cdot \mathbf{L} \right) \Psi_+ + (E + 2m - \Delta)(E - \Sigma)\Psi_+ \quad (2)$$

$$p^2\Psi_- = \frac{\Sigma'}{E - \Sigma} \left(\frac{\partial}{\partial r} - \frac{1}{r} \boldsymbol{\sigma} \cdot \mathbf{L} \right) \Psi_- + (E + 2m - \Delta)(E - \Sigma)\Psi_- , \quad (3)$$

where $\Psi_{\pm} = [(1 \pm \beta)/2]\Psi$ are the upper and lower components and the primes denote derivatives with respect to r . The $\boldsymbol{\sigma} \cdot \mathbf{L}$ term in (3) is the pseudospin-orbit term.

We solve the Dirac equation with the Hamiltonian (1) with scalar and vector potentials with Woods-Saxon shape: $U(r) = U_0/(1 + \exp[(r - R)/a])$. We fitted the parameters of this potential to the neutron spectra of ²⁰⁸Pb, obtaining the values $R = 7$ fm, $\Delta_0 = 650$ MeV, $\Sigma_0 = -66$ MeV and $a = 0.6$ fm. We then varied the diffuseness, radius and Σ_0 separately, keeping all the other parameters fixed and showed that the splittings of the pseudospin doublets vary notably, namely they decrease as $|\Sigma_0|$ decreases and diffuseness increases, sometimes reversing sign. The details can be found in ref. ⁷.

The effect of the pseudospin-orbit term on pseudospin energy splittings can be assessed by writing (3) as a Schroedinger-like equation, dividing it by an energy- and r -dependent effective mass $m^* = (E + 2m - \Delta)/2$ and taking the expectation value of each term relative to Ψ_- :

$$\left\langle \frac{p^2}{2m^*} \right\rangle + \langle V_{\text{PSO}} \rangle + \langle V_{\text{D}} \rangle + \langle \Sigma \rangle = E, \quad (4)$$

where

$$\left\langle \frac{p^2}{2m^*} \right\rangle = \int \Psi_-^\dagger \frac{p^2}{2m^*} \Psi_- d^3\mathbf{r} / \int \Psi_-^\dagger \Psi_- d^3\mathbf{r}, \quad (5)$$

$$\langle V_{\text{PSO}} \rangle = P \int \Psi_-^\dagger \frac{1}{2m^*} \frac{\Sigma'}{E - \Sigma} \frac{1}{r} \boldsymbol{\sigma} \cdot \mathbf{L} \Psi_- d^3\mathbf{r} / \int \Psi_-^\dagger \Psi_- d^3\mathbf{r} \quad (6)$$

$$\langle V_D \rangle = -P \int \Psi_-^\dagger \frac{1}{2m^*} \frac{\Sigma'}{E - \Sigma} \frac{\partial \Psi_-}{\partial r} d^3\mathbf{r} / \int \Psi_-^\dagger \Psi_- d^3\mathbf{r}, \quad (7)$$

$$\langle \Sigma \rangle = \int \Psi_-^\dagger \Sigma \Psi_- d^3\mathbf{r} / \int \Psi_-^\dagger \Psi_- d^3\mathbf{r}, \quad (8)$$

where ‘P’ denotes the principal value of the integral. These terms can be identified, respectively, as a kinetic term, a pseudospin-orbit term, a potential term related to what is sometimes called Darwin term and the mean value of the Σ potential with respect to the lower component, Ψ_- .




Figure 1. Differences for the energy terms in (4) for $(1i_{11/2}, 2g_{9/2})$, $(2f_{5/2}, 3p_{3/2})$ and $(1h_{9/2}, 2f_{7/2})$ pseudospin partners.




Figure 2. Effect of Coulomb and ρ potentials V_c and V_ρ in the splitting of the $(2d_{5/2}, 1g_{7/2})$ pseudospin partners in a Sn isotopic chain.

In Fig. 1 are plotted the differences of the terms in Eq. (4) between each member of the three topmost pseudospin partners in ^{208}Pb , $(1i_{11/2}, 2g_{9/2})$, $(2f_{5/2}, 3p_{3/2})$ and $(1h_{9/2}, 2f_{7/2})$. One sees that the contribution of V_{PSO} for the pseudospin energy splittings is larger than the splittings themselves and has the opposite sign of the kinetic and $\langle \Sigma \rangle$ terms. For all doublets there is a significant cancellation of $\langle V_{\text{PSO}} \rangle$ with the kinetic and $\langle \Sigma \rangle$ terms. A similar analysis was made in ref. ⁶, concluding that the pseudospin-orbit potential is non-perturbative. We also found that there is a clear correlation between $\langle V_{\text{PSO}} \rangle$ and the pseudospin energy splitting when the diffusivity and the depth Σ_0 are varied ⁷.

All these findings point to the conclusion that pseudospin symmetry has a dynamical nature, since it is very much dependent on the shape of the Σ nuclear mean field potential and results from a strong cancellation among the several contributing terms to the energy splittings.

This pseudospin dependence on the shape of the Σ potential allows us to explain why the proton and neutron levels have different pseudospin splittings⁸.

In RMF models, the ρ meson interaction modifies the vector potential to $V = V_\omega + V_\rho = V_\omega \pm \frac{g_\rho}{2}\rho_0$, where the plus sign refers to protons and the minus sign refers to neutrons, and ρ_0 is the time component of the ρ field which is proportional to $Z - N$. The magnitude of the ρ potential is around 4-8 MeV. For heavy nuclei, $\rho_0 < 0$ and therefore $V_p < V_n$ which implies $|\Sigma_n| < |\Sigma_p|$. This means that, for neutron-rich nuclei, the pseudospin symmetry is favored for neutrons. Notice, however, that the Coulomb potential, which has the opposite sign of V_ρ for protons, also affects proton spectra. Both of these effects can be seen in Figure 2, where is plotted the effect of V_ρ and the Coulomb potential on the energy splitting of the $(2d_{5/2}, 1g_{7/2})$ pseudospin partners along the Sn isotopic chain from $A = 130$ to $A = 170$. We see that the potential V_ρ gives the main contribution for the observed isospin asymmetry.

Acknowledgments

We acknowledge financial support from FCT (POCTI), Portugal, and from CNPq/ICCTI Brazilian-Portuguese scientific exchange program.

References

1. K. T. Hecht and A. Adler, Nucl. Phys. **A137**, 129 (1969)
2. A. Arima, M. Harvey, and K. Shimizu, Phys. Lett. **B30**, 517 (1969)
3. J. N. Ginocchio, Phys. Rev. Lett. **78**, 436 (1997); *ibid*, Phys. Rept. **315**, 231 (1999)
4. J. N. Ginocchio and A. Leviatan, Phys. Lett. **B245**, 1 (1998) *ibid*, Phys. Rev. Lett. **82**, 4599 (1999)
5. J. Meng, K. Sugawara-Tanabe, S. Yamaji, P. Ring, and A. Arima, Phys. Rev. **C58**, R628 (1998)
6. S. Marcos, M. López-Quelle, R. Niembro, L. N. Savushkin, and P. Bernardos, Phys. Lett. **B513**, 30 (2001)
7. P. Alberto, M. Fiolhais, M. Malheiro, A. Delfino, and M. Chiapparini, Phys. Rev. Lett. **86**, 5015 (2001); *ibid*, Phys. Rev. C **65**, 034307 (2002)
8. J. Meng and I. Tanihata, Nucl. Phys. **A650**, 176 (1999)